

**Solutions to homework set 8 — APPM5440 — Fall 2012**

**3.6:** Set  $X = C([-a, a])$  and define on  $X$  the operator

$$[Fu](x) = \frac{1}{\pi} \int_{-a}^a \frac{1}{1 + (x - y)^2} u(y) dy + 1.$$

Then the given equation can be formulated as a fixed point problem  $u = F(u)$ . We find that

$$\|F(u) - F(v)\| = \sup_{x \in [-a, a]} \left| \frac{1}{\pi} \int_{-a}^a \frac{1}{1 + (x - y)^2} (u(y) - v(y)) dy \right| \leq \dots \leq \frac{2 \arctan(a)}{\pi} \|u - v\|$$

(you should be able to fill in the missing steps). The contraction mapping principle now proves that the equation has a unique solution in  $X$ .

To prove that the solution is positive, note that if  $u(x) \geq 0$  for all  $x$ , then  $[Tu](x) \geq 0$  for all  $x$ . Combine this fact with the fact that  $f$  is non-negative, and that

$$u = (I - T)^{-1} f = \sum_{n=0}^{\infty} T^n f.$$

For the case  $a = \infty$ , note that uniqueness cannot hold. If  $u$  is a solution, then so is any shift of  $u$ .

**3.7:** Set  $X = \{u \in C(I) : u(0) = u(1) = 0\}$ . Convolve given BVP with the Green's function  $g(x, y)$  (defined by eqn (3.22)) and obtain

$$(1) \quad u + \lambda T u = h,$$

where  $T$  is the operator on  $X$  given by

$$[Tu](x) = \int_0^1 g(x, y) \sin(u(y)) dy,$$

and

$$h(x) = \int_0^1 g(x, y) f(y) dy.$$

Note that  $Tf \in X$ , so (1) is an equation on  $X$ . Now

$$\|Tu - Tv\| \leq \sup_x \int_0^1 |g(x, y)| |\sin(u(y)) - \sin(v(y))| dy.$$

Use that

$$|\sin(u(y)) - \sin(v(y))| \leq |u(y) - v(y)| \leq \|u - v\|$$

to obtain

$$\|Tu - Tv\| \leq \beta \|u - v\|,$$

where

$$\beta = \sup_x \int_0^1 |g(x, y)| dy.$$

We see that if  $\lambda < 1/\beta$ , equation (1) has a unique solution in  $X$  (by the contraction mapping theorem).

**Problem 4:** Consider the integral equation

$$(*) \quad u(x) = \pi^2 \sin(x) + \frac{3}{2} \int_0^{\cos(x)} |x - y| u(y) dy.$$

Prove that (\*) has a unique solution in  $C([0, 1])$ .

**Solution:** Set  $X = C([0, 1])$  and define the operator  $T$  on  $X$  by

$$[Tu](x) = \frac{3}{2} \int_0^{\cos(x)} |x - y| u(y) dy.$$

We find that

$$\begin{aligned} \|Tu - Tv\| &\leq \sup_x \frac{3}{2} \int_0^{\cos(x)} |x - y| |u(y) - v(y)| dy \\ &\leq \sup_x \frac{3}{2} \int_0^1 |x - y| |u(y) - v(y)| dy \\ &\leq \sup_x \frac{3}{2} \int_0^1 |x - y| dy \|u - v\|. \end{aligned}$$

Now

$$\sup_x \int_0^1 |x - y| dy = \sup_x \left( \frac{1}{2}x^2 + \frac{1}{2}(1 - x)^2 \right) = \frac{1}{2},$$

so

$$\|Tu - Tv\| \leq \frac{3}{4} \|u - v\|.$$

Since  $T$  is a contraction, the equation

$$(I - T)u = f$$

has a unique solution for every  $f$ . (In particular, for  $f(x) = \pi^2 \sin(x)$ .)

**Problem 1:** Let  $X$  be a set with infinitely many members. We define a collection  $\mathcal{T}$  of subsets of  $X$  by saying that a set  $\Omega \in \mathcal{T}$  if either  $\Omega^c = X \setminus \Omega$  is finite, or if  $\Omega$  is the empty set. Verify that  $\mathcal{T}$  is a topology on  $X$ . This topology is called the “co-finite” topology on  $X$ . Describe the closed sets.

**Solution:** The verification should be straight-forward. The closed sets are the finite sets, and the entire set.

**Problem 2:** Let  $X$  denote a finite set, and let  $\mathcal{T}$  be a metrizable topology on  $X$ . Prove that  $\mathcal{T}$  is the discrete topology on  $X$ .

**Solution:** Enumerate the points in  $X$  so that  $X = \{x_n\}_{n=1}^N$ . Set

$$\varepsilon_n = \min\{d(x_n, x_m) : m \neq n\}, \quad \text{for } n = 1, 2, \dots, N.$$

Since the minimum is taken over a finite set of positive numbers,  $\varepsilon_n > 0$ , and it follows that the set  $B_{\varepsilon_n/2}(x_n) = \{x_n\}$ , must be open. Since any union of open set must itself be open, it follows that all subsets of  $X$  are open.

**Problem 3:** Consider the set  $X = \{a, b, c\}$ , and the collection of subsets  $\mathcal{T} = \{\emptyset, \{a\}, \{a, b\}, \{a, c\}, \{a, b, c\}\}$ . Is  $\mathcal{T}$  a topology? Is  $\mathcal{T}$  a metrizable topology?

**Solution:** Yes it is a topology (and union or intersection of the given sets is itself a member of the set.

No, it cannot be metrizable. If it were, then the argument given in Problem 2 would demonstrate that  $\{b\}$  would have to be an open set, and it is not.