

Homework set 9 — APPM5450, Spring 2006

From the textbook: 11.18, 11.13, 11.15, 11.16.

In 11.16, you're free to assume that f is smooth (or that $f \in \mathcal{S}(\mathbb{R}^3)$), if you like.

If you had difficulties with Midterm 2, then you may find the following review questions useful:

What does it mean for $\varphi_n \rightarrow \varphi$ in \mathcal{S} ?

Make absolutely sure that you understand problems like 11.4, 11.10a.

Prove that if $\varphi_n \rightarrow \varphi$ in \mathcal{S} , then $x\varphi_n(x) \rightarrow x\varphi(x)$ and $\partial\varphi_n \rightarrow \partial\varphi$ in \mathcal{S} .

Let T be a linear map from \mathcal{S} to \mathbb{R} . What does it mean for T to be continuous? Prove that if there exists a finite C and a finite N such that $|T(\varphi)| \leq C \sum_{|\alpha|, n \leq N} \|\varphi\|_{\alpha, n}$, then T is continuous.

Let $T \in \mathcal{S}^*(\mathbb{R}^d)$, and let α be a multi-index. Define $x^\alpha T$. Prove that what you define is a tempered distribution.

Prove that $n^2 \sin(nx) \rightarrow 0$ in \mathcal{S}^* .

Is the Schwartz space dense in \mathcal{S}^* ?

Prove that $\sup_x |x^\beta \partial^\alpha \varphi(x)| < \infty \forall \alpha, \beta \Leftrightarrow \sup_x |(1+|x|^2)^{k/2} \partial^\alpha \varphi(x)| < \infty \forall \alpha, k$.

Assume that $\int |f|^2 < \infty$, set $\langle T, \varphi \rangle = \int f \varphi$. Prove that $T \in \mathcal{S}^*$.

Let H be a function such that $H(x) = 1$ if $x \geq 0$, zero otherwise. Prove that $H \in \mathcal{S}^*$. Calculate H' . Let H_R denote the function that is 1 when $0 \leq x \leq R$ and zero otherwise. Prove that $H_R \rightarrow H$ in \mathcal{S}^* as $R \rightarrow \infty$.

Let ψ be a Schwartz function such that $\int \psi = 0$. Set $\varphi_n(x) = n\psi(nx)$. Does φ_n converge in \mathcal{S} ? Does φ_n converge in \mathcal{S}^* ?

Prove that $\text{PV}(1/x)$ is a continuous functional on \mathcal{S} .

What is the distributional derivative of $\text{PV}(1/x)$?