

## Homework set 4 — APPM5450, Spring 2007

From the textbook: 8.15.

**Problem 1:** Consider the Hilbert space  $H = l^2(\mathbb{N})$ ; let  $e_n$  denote the canonical basis vectors. Which of the following sequences converge weakly? Which have convergent subsequences?

- (a)  $x_n = n e_n$ .
- (b)  $y_n = n^{-1/2} \sum_{j=1}^n e_j$ .
- (c)  $x_n = e_n + e_m$  where  $m = 1 + \text{mod}(n, 2)$ .

**Problem 2:** Consider the Hilbert space  $H = L^2([-\pi, \pi])$ , and the sequence of functions  $\varphi_n(x) = x^2 \sin(nx)$ . Does  $(\varphi_n)_{n=1}^\infty$  converge strongly in  $H$ ? Does  $(\varphi_n)_{n=1}^\infty$  converge weakly in  $H$ ? If you answer yes to either question, specify the limit.

**Problem 3:** Let  $A$  denote a self-adjoint operator on a Hilbert space  $H$ . Let  $u$  denote an element of  $H$  and set  $u_n = e^{i n A} u$ . Prove that  $(u_n)_{n=1}^\infty$  has a weakly convergent subsequence.

**Problem 4:** Let  $H_1$  and  $H_2$  be two Hilbert spaces. Let  $U : H_1 \rightarrow H_2$  be a unitary operator, and let  $A_1 \in \mathcal{B}(H_1)$  be a self-adjoint operator. Define the operator  $A_2 \in \mathcal{B}(H_2)$  by  $A_2 = U A_1 U^{-1}$ . Prove that  $A_2$  is self-adjoint.

**Problem 5 (optional):** Consider the Hilbert space  $H = L^2([-\pi, \pi])$ , and let  $\mathcal{P}$  denote the set of trigonometric polynomials (which is dense in  $H$ ). For  $u \in \mathcal{P}$ , let  $A$  denote the operator  $Au = 100u + 18u'' + u''''$ . Prove that

$$\sup_{u \in \mathcal{P}, \|u\|=1} \langle Au, u \rangle = \infty.$$

Conclude that  $A$  cannot be extended to a bounded linear operator on  $H$ . Prove that for  $u, v \in \mathcal{P}$ ,  $\langle Au, v \rangle = \langle u, Av \rangle$ . Determine

$$\inf_{u \in \mathcal{P}, \|u\|=1} \langle Au, u \rangle.$$

Prove that

$$\langle u, v \rangle_A = \langle Au, v \rangle$$

is a bilinear form on  $\mathcal{P}$ . Prove that on  $\mathcal{P}$ , the norm  $\|\cdot\|_A$  induced by  $\langle \cdot, \cdot \rangle_A$  is equivalent to the norm

$$\|u\|_{H^2(\mathbb{T})} = \sqrt{\|u\|_{L^2(\mathbb{T})}^2 + \|u''\|_{L^2(\mathbb{T})}^2}.$$

Conclude that the closure of  $\mathcal{P}$  under the norm  $\|\cdot\|_A$  is the space  $H^2(\mathbb{T})$  (as defined in Section 7.2).